

67 GTG TCA ACC TGG TTG ATA CGT AGT GGT GAA CCT GTG CAA CAC CGC ACT GAA TTC CCG TTC
 127 ATC GCA TTC TTA ACG ACA GAG AGA ACA ATG TGT ACA GGT TCA CTA GTC TCA ACG AGA GCA
 187 GTA CTC ACA GCT GGT CAT TGT GTT TGC TCA CCA TTG CCA GTG ATT CGG gtgagatcgactg
 250 aaacacattgttgtctcaatgtaacgattgatttcacagacagtggttgcgtgtgctgctggtttgtgtttgtgtatgca
 329 gtgggttgcatagatttttaattgtctatataacttgggtcttatttcag/gtt TCA TTT CTC ACA CTG AGG AAT
 400 GGC GAC CAA CAA GGC ATC CAT CAC CAA CCG TCT GGA GTT AAG GTG GCA CCA GGA TAC ATG
 460 CCC TCT TGT ATG TCG GCA CGA CAG AGG AGA CCA ATC GCA CAG ACA CTC AGT GGA TTC GAT
 520 ATT GCA ATT GTA ATG CTG GCT CAA ATG GTC AAC TTA CAG AGT GGA ATC AGA GTG ATC AGT
 580 CTG CCA CAG CCA TCG GAT ATC CCG CCA CCT GGA ACT GGT GTT TTC ATT GTT GGT TAT GGA
 640 AGG GAT GAT AAC GAC CGT GAT CCG TCA CGT AAG AAT GGT GGA ATA TTG AAG AAA gtgagat
 701 gttggggaataaacgacatgactcagtcagtcagtcagatgtcagtcagttattctgtgtgtctgtgtatctgtttgtgt
 780 gtctgtctgtctacctgatccggttgttgtattgggtcagagcccttgataataacaactgtgtttggatgactttgtgac
 859 agttcagtagcagagtgatttccatctcggtcattgtgttggtaggtgaggtgacgtgatgtgaggtgagttgaggga
 938 gatggatgggatggaatgtgatgtgatgggatgattgagaccacttggaggagagaagactcatgaaatatctatgca
 1017 aacgatggaagtgtgttgtgtacatgaagttaggggtcaatgtgtttgagtatgtgtttggagagtggtagatggaga
 1096 gtgacttgatcgtcgaatatagtgcacatgtgattgtatgtggactattgttgtgtgggtagtgtgaaggggtggatatt
 1175 gtgccagttgatattttcgaattcacttgtgtgttttgttttgttccgttgtgttttgttctttctcatctgtactgt
 1254 acgttgttgtactgtactgttgtgtgttgtttgtttgcaccacag GGT CGA GCG ACT ATA ATG GAA TGC
 1325 CGA CAT GCG ACC AAT GGC AAT CCT ATA TGT GTG AAA GCA GGT CAG AAT TTC GGA CAG TTA
 1385 CCC GCT CCA GGT GAC AGT GGT GGA CCT CTC CTC CCA TCC CTT CAA GGT CCA GTA CTC GGT
 1445 GTC GTA TCA CAT GGT GTC ACA CTG CCA AAC CTT CCC GAT ATC ATT GTC GAG TAT GCC AGT
 1505 GTG GCT AGA ATG TTG GAT TTT GTA CGC TCC AAT ATT TGA

Fig. 1. Consensus sequence of 3 PCR clones of *Schistosoma mansoni* cercarial elastase gene. Nucleotide numbers correspond to the open reading frame of the cDNA sequence reported by Newport *et al.* (1988). The 2 introns are shown in lower case text. Splice signal and branch formation sequences are shown in underlined text.

BEST AVAILABLE COPY

1002044-121004

68
 70
 0
 18
 96
 50
 30 - 687 X - Cur
 Full length
 67 -
 106
 2
 3

Figure 2. - Production of Construct pGEXCEL





Figure 3 - SDS-PAGE Showing Purification of the Recombinant Protein Sm30-GST




- M = Protein Mid-Range Molecular Weight Markers**
1 = Total proteins from uninduced *E. coli* carrying plasmid pGEXCEL
2 = Total proteins from IPTG-induced *E. coli* carrying pGEXCEL, grown at 30°C
3 = Soluble lysate of IPTG-induced *E. coli* as in lane 2
4 = Fraction containing Sm30-GST following ion-exchange chromatography
5 = Fraction containing Sm30-GST following gel filtration

The recombinant protein Sm30-GST is indicated by "←"

Figure 4:

Western immunoblot showing antibody reactivity of serum from rabbit immunized with Sm30-GST.




Antigens in lanes:

1. Horse albumin
2. Recombinant fusion protein Sm30-GST
3. " " " Sm30Ex1-GST
4. " " " Sm30Ex2-GST →
5. " " " Sm30Ex3-GST
6. Recombinant Sj26GST
7. *S. mansoni* cercarial transformation fluid (CTF)

Figure 5

Reactivity of sera from immunized mice against *S. mansoni* cercarial antigens.



A = sera from 6 mice immunized with *S. mansoni* CTF

B = sera from 6 mice immunized with Sm30-GST

1 = serum from rabbit BR67


2 = normal rabbit serum

3 = larval antigens stained with protogold

M = molecular weight standards

Figure 6


Reactivity of sera from immunized mice against *S. haematobium* cercarial antigens.



A = sera from 6 mice immunized with *S. mansoni* CTF
 B = sera from 6 mice immunized with Sm30-GST
 1 = serum from rabbit BR67
 2 = normal rabbit serum
 3 = larval antigens stained with protogold
 M = molecular weight standards

Figure 7:

Cross-reactivity of antibodies raised against *S. haematobium* 27kDa larval protease against antigens of *S. mansoni* and *S. margrebowiei*.




Antigens in lanes:

1. *S. mansoni* cercarial antigens
 2. *S. haematobium* cercarial antigens
 3. *S. margrebowiei* cercarial antigens
- M = molecular weight standards (kDa)

Rabbit antiserum used to probe antigens in lanes 1-3 was raised against the 27kDa *S. haematobium* larval protease using the same methods as for BR67, the anti-*S. mansoni* 27kDa larval protease antiserum.

Figure 8:

Antibody reactivity of mice immunized with recombinant fusion proteins against *S. mansoni* cercarial transformation fluid (CTF).



Group	Immunogen
A	Alum and PBS alone (control mice).
B	CTF.
C	Recombinant Sj26.
D	Recombinant Sm30-GST.
E	Recombinant Sm30Ex1-GST
F	Recombinant Sm30Ex2-GST
G	Recombinant Sm30Ex3-GST

Control Sera

E = Rabbit serum BR67 (anti-27kDa *S. mansoni* larval protease)

G = Rabbit serum 1093X (anti-*S. japonicum* GST)

N = Normal rabbit serum

M = Molecular weight markers.

M =

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.